



ROI OF SOYBEAN FOLIAR INSECTICIDES

ACTION TRIAL PROTOCOL

In partnership with **UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN**

PROTOCOL AUTHORS – Nick Seiter, University of Illinois

Reviewed by the ISA OFTN Team

OBJECTIVE – Determine how often soybean insecticide foliar applications provide a positive return in protected yield and which insect pests are responsible when they do.

- Number of locations: 4 on-farm sites
- Geographic scope: Statewide

TREATMENTS – Comparison of commercial foliar insecticide treatment to control treatment without this insecticide applied.

Insecticide Treatment:

- Foliar insecticide application at R3

Control Requirements:

- Except for foliar insecticide being compared, all other management should be the same as the plots with insecticide treatment.

Treatment Rationale:

- The specific treatment is flexible depending on the interest of the farmer. For instance, we could test the addition of a pyrethroid insecticide to a planned fungicide application, compare a premix insecticide to a single active ingredient, or compare fungicide + insecticide seed treatment with fungicide-only.

PLOT LAYOUT – Flexible depending on equipment and field dimensions.

- Minimum field size of 40 acres.
- Trials will consist of at least 4 replications of both a control and foliar insecticide treatment (total of 8 plots), each a minimum of 60 feet wide.
- Width of plots can be adapted to farmer's equipment.
 - Must be wide enough to measure yield using farmer's harvest equipment.
 - Minimum width is the larger of either 60 ft or twice the width of the combine header (to avoid edge effect).

ROI OF SOYBEAN FOLIAR INSECTICIDES

ACTION TRIAL PROTOCOL

EQUIPMENT NEEDED

- Appropriate equipment to apply treatment
 - Foliar spray equipment
 - Drone application service within range and able to be contracted
- Combine with a calibrated yield monitor

FIELD WORK AND MEASUREMENTS COLLECTED

- Agronomic management information - *Farmer provided*
 - Previous 3-5 years of basic management information preferred
 - More detailed current year's management information
- Field applications - *Farmer responsible*
- Baseline soil fertility sampling & analysis - *ISA responsible, spring*
- Insect counts and injury assessments, 4-8 times throughout the season - *performed by the Seiter laboratory*
- Yield monitor data - *Farmer provided*

FOLIAR APPLICATION PROCESS

- Insecticide will be applied to the field in replicated strips according to a plot map coordinated with ISA. This map can be imported as a shapefile into most spray drones.
- An application plan (whether applied using a ground rig or drone), including who will make the application (farmer, custom applicator, etc.) should be approved prior to enrollment in the protocol.
- The insecticide used will be chosen in consultation with the farmer based on interest, previous use, etc., but we have typically used a pyrethroid like lambda-cyhalothrin or bifenthrin.

ISA OFTN CONTACT INFO

Deanna Burkhart

Agronomic Services Manager

deanna.burkhart@ilsoy.org

309.307.9366

Illinois Soybean Association

Main Office

1108 Trinity Lane

Bloomington, IL 61704