

Added on: July 6, 2023 at 2:15 pm

**Select Your Form**

Soy Envoys/ISA Agronomy Team/Others

**Name**

Leonardo Rocha

**Email**

leonardo.rocha@siu.edu

**Date**

July 6, 2023

**County**

White

**Region/State**

Region 6

**Field Photo Upload 1**



**Field Photo Upload 2**



### Field Photo Upload 3



### Field Photo Upload Caption 1

Double crop soybean emerging

### Field Photo Upload Caption 2

Double crop soybean at VC stage

### Field Photo Upload Caption 3

Japanese beetles causing damage in soybean

### Which of the following best describes current conditions in this county?

Mildly Wet (soil is wetter than normal, local vegetation is healthy)

If conditions are on the dry end, which of the following US Drought Monitor categories best fit current conditions. To better judge the fit, see explanation of USDM categories here:

<https://droughtmonitor.unl.edu/About/AbouttheData/DroughtClassification.aspx>

Near Normal (Dnada)

### Quick synopsis of conditions that will appear in the main feed

The recent precipitation events left the soil with high moisture levels, helping double cropping soybean fields to emerge. This particular field was planted about two weeks ago, under drought and without access

irrigation, but plants are emerging with high vigor and uniform stand. With more rain on the radar, this particular field has good yield potential at this point.

### **Weather**

Cloudy, temperatures in the mid 80s

### **Soybean Growth Stage**

Some fields reaching R4, R1-R3 in general

### **Corn Growth Stage**

VT-R1

### **Insects**

Some Japanese beetle damage, but probably not even close to damage thresholds

### **Weeds**

Hophornbeam copperleaf widespread in some areas, patches of nightshade

Marestail in double crop fields